

- Abedrabbo, S. (1994). Control of the little fire ant, *Wasmannia auropunctata*, on Santa Fe islands in the Galapagos islands. In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 219–227). Boulder CO: Westview Press.
- Abensperg-Traun, M. (1992). Biomass of surface-foraging ants (Formicidae) in four bushland habitats in the wheatbelt of Western Australia. *J. R. Soc. West. Aust.*, 75, 25–32.
- Abensperg-Traun, M., Smith, G. T., Arnold, G. W. & Steven, D. E. (1996). The effects of habitat fragmentation and livestock-grazing on animal communitites in remnants of gimlet *Eucalyptus salubris* woodland in the Western Australian wheatbelt. I. Arthropods. *J. Appl. Ecol.* 33, 1281–1301.
- Acosta, F. J., Serrano, J. M. & Zorrilla, J. M. (1988). La densidad como condicionante de la autoorganizacion espacial en comunidades de hormigas. *Rev. Ecol. Biol. Sol.*, 25, 115–127.
- Acosta, F. J., Zorrilla, J. M., López, F. & Serrano, J. M. (1992). Progressive coupling of ant community topography with environment geomorphology in post perturbed slopes. *Acta Entomol. Bohemslav.*, 89, 157–167.
- Adams, E. S. & Tschinkel, W. R. (1995). Spatial dynamics of colony interactions in young populations of the fire ant *Solenopsis invicta*. *Oecologia*, 102, 156–163.
- Adams, E. S. & Tschinkel, W. R. (1995). Density-dependent competition in fire ants: effects on colony survivorship and size variation. *J. Anim. Ecol.*, 64, 315–324.
- Addison, P. & Samways, M. J. (2000). A survey of ants (Hymenoptera: Formicidae) that forage in vineyards in the Western Cape Province, South Africa. *Afr. Entomol.*, 8, 251–260.
- Aktaç, N. (1978). Seasonal changes of the ant nests number. *Biyol. derg.*, 28, 53–65.
- Albrecht, M. & Gotelli, N. J. (2001). Spatial and temporal niche partitioning in grassland ants. *Oecologia*, 126, 134–141.

Alvarado, M. & Gallé, L. (1999). Diversity and structure of ant assemblages in forest habitats in the South Hungarian Plain. *Soil Zoology in Central Europe. Proc. 5th Centr. Europ. Workshop Soil Zool.*, 1–6.

Alvarado, M. (2000). Habitat correlates of ant assemblages in different forests of the south Pannonian plain. *Tiscia*, 32, 35–42.

Andersen, A. N. (1983). A brief survey of ants in Glenadale National Park, with particular reference to seed-harvesting. *Vict. Nat.*, 100, 233–237.

Andersen, A. N. & McKaige, M. E. (1987). Ant communities at Rotamah island, Victoria, with particular reference to disturbance and *Rhytidoponera tasmaniensis*. *Proc. R. Soc. Vict.*, 99, 141–146.

Andersen, A. N. (1988). Immediate and longer-term effects of fire on seed predation by ants in sclerophyllous vegetation in south-eastern Australia. *Aust. J. Ecol.*, 13, 285–293.

Andersen, A. N. (1990). The use of ant communities to evaluate change in Australian terrestrial ecosystems: a review and a recipe. *Proc. Ecol. Soc. Aust.*, 16, 347–357.

Andersen, A. N. (1991). Responses of ground-foraging ant communities to three experimental fire regimes in a savanna forest of tropical Australia. *Biotropica*, 23, 575–585.

Andersen, A. N. & Burbidge, A. H. (1991). The ants of a vine thicket near Broome: a comparison with the northwest Kimberley. *J. R. Soc. West. Aust.*, 73, 79–82.

Andersen, A. N. & Majer, J. D. (1991). The structure and biogeography of rainforest ant communities in the Kimberley region of northwestern Australia. In McKenzie, N.L., Johnston, R. B., Kendrick, P. G. (Eds.), *Kimberley rainforests* (pp. 333–346). Chipping Norton, NSW: Surrey Beatty & Sons.

Andersen, A. N., Myers, B. A. & Buckingham, K. M. (1991). The ant fauna of a mallee outlier near Melton, Victoria. *Proc. R. Soc. Vict.*, 103, 1–6.

- Andersen, A. N. (1992). Regulation of "momentary" diversity by dominant species in exceptionally rich ant communities of the Australian seasonal tropics. *Am. Nat.*, 140, 401–420.
- Andersen, A. N. (1993). Ant communities in the Gulf region of Australia's semi-arid tropics: Species composition, patterns of organisation, and biogeography. *Aust. J. Zool.*, 41, 399–414.
- Andersen, A. N. & Patel, A. D. (1994). Meat ants as dominant members of Australian ant communities: an experimental test of their influence on the foraging success and forager abundance of other species. *Oecologia*, 98, 15–24.
- Andersen, A. N. & Reichel, H. (1994). The ant (Hymenoptera: Formicidae) fauna of Holmes Jungle, a rainforest patch in the seasonal tropics of Australia's Northern Territory. *J. Aust. ent. Soc.*, 33, 153–158.
- Andersen, A. N. & Spain, A. V. (1996). The ant fauna of the Bowen Basin, in the semi-arid tropics of Central Queensland (Hymenoptera: Formicidae). *Aust. J. Entomol.*, 35, 213–221.
- Andersen, A. N. (1997). Functional groups and patterns of organization in North American ant communities: a comparison with Australia. *J. Biogeogr.*, 24, 433–460.
- Andersen, A. N. (1997). Ants as indicators of ecosystem restoration following mining: a functional group approach. In Hale, P. & Lamb, D. (Eds.), *Conservation outside nature reserves*. (pp. 319–325). Queensland.
- Andersen, A. N., Azcarate, F. M. & Cowie, I. D. (2000). Seed selection by an exceptionally rich community of harvester ants in the Australian seasonal tropics. *J. Anim. Ecol.*, 69, 975–984.
- Andersen, A. N. & Müller, W. J. (2000). Arthropod responses to experimental fire regimes in an Australian tropical savannah: ordinal-level analysis. *Austral Ecol.*, 25, 199–209.

- Andersen, A. N. (2003). Ant biodiversity in arid Australia: productivity, species richness and community organisation. *Rec. South Aust. Mus. Monogr. Ser.*, 7, 79–92.
- Andersen, A. N., Hoffmann, B. D. & Somes, J. (2003). Ants as indicators of minesite restoration: community recovery at one of eight rehabilitation sites in central Queensland. *Ecol. Manag. Restor.*, 4, S12–S19.
- Andrew, N., Rodgerson, L. & York, A. (2000). Frequent fuel-reduction burning: the role of logs and associated leaf litter in the conservation of ant biodiversity. *Austral Ecol.*, 25, 99–107.
- Armbrecht, I. & Ulloa-Chacon, P. (1999). Rareza y diversidad de hormigas en fragmentos de bosque seco Colombianos y sus matrices. *Biotropica*, 31, 646–653.
- Armbrecht, I., Jimenez, E., Alvarez, G., Ulloa-Chacon, P. & Armbrecht, H. (2001). An ant mosaic in the Colombian rain forest of Choco (Hymenoptera: Formicidae). *Sociobiology*, 37, 491–509.
- Armbrecht, I., Perfecto, I. & Vandermeer, J. (2004). Enigmatic biodiversity correlations: ant diversity responds to diverse resources. *Science*, 304, 284–286.
- Arnold, G. W., Abensperg-Traun, M., Hobbs, R. J., Steven, D. E., Atkins, L., Viveen, J. J. & Gutter, D. M. (1999). Recovery of shrubland communities on abandoned farmland in south-western Australia: Soils, plants, birds and arthropods. *Pac. Conserv. Biol.*, 5, 163–178.
- Assing, V. (1986). Distribution, densities and activity patterns of the ants (Hymenoptera: Formicidae) of Calluna Heathlands in Northwestern Germany. *Entomol. Gener.*, 11, 183–190.
- Assing, V. (1989). Die Ameisenfauna (Hym.: Formicidae) nordwestdeutscher *Calluna*-Heiden. *Drosera*, 89, 49–62.
- Azcarate, F. M. & Peco, B. (2003). Spatial patterns of seed predation by harvester ants

- (*Messor* Forel) in Mediterranean grassland and scrubland. *Insect. Soc.*, 50, 120–126.
- Baroni Urbani, C. & Aktaç, N. (1981). The competition for food and circadian succession in the ant fauna of a representative Anatolian semi-steppic environment. *Mitt. Schweiz. Entomol. Ges.*, 54, 33–56.
- Baroni Urbani, C. (1987). Comparative feeding strategies in two harvesting ants. In Eder, J. & Rembold, H. (Eds.), *Chemistry and biology of social insects* (pp. 509–510). München: Verlag J. Peperny.
- Basedow, T. (1993). Predatory arthropods in cabbage terraces under different conditions in the Cordillera region of Luzon, Philippines. *Bull. Entomol. Res.*, 83, 313–319.
- Behr, D. & Cölln, K. (1993). Zur Ameisenfauna (Hymenoptera, Formicidae) von Gönnersdorf (Kr. Daun). *Dendrocopos*, 20, 148–160.
- Behr, D. & Cölln, K. (1994). Ameisen (Formicidae) einer Eifellandschaft mit Untersuchungen zur Pflege von Kalkmagerrasen sowie einer vorläufigen Gesamtarteliste für den Eifelraum. *Dendrocopos*, 21, 121–146.
- Bernstein, R. A. & Gobbel, M. (1979). Partitioning of space in communities of ants. *J. Anim. Ecol.*, 48, 931–942.
- Bestelmeyer, B. T. & Wiens, J. A. (1996). The effects of land use on the structure of ground-foraging ant communities in the Argentine chaco. *Ecol. Appl.*, 6, 1225–1240.
- Bestelmeyer, B. T. (1997). Stress tolerance in some Chacoan dolichoderine ants: implications for community organization and distribution. *J. Arid. Environ.*, 35, 297–310.
- Bestelmeyer, B. T. & Schooley, R. L. (1999). The ants of the southern Sonoran desert: community structure and the role of trees. *Biodiv. Conserv.*, 8, 643–657.
- Bestelmeyer, B. T. (2000). The trade-off between thermal tolerance and behavioural dominance in a subtropical South American ant community. *J. Anim. Ecol.*, 69, 998–1009.

- Bestelmeyer, B. T. & Wiens, J. A. (2001). Ant biodiversity in semiarid landscape mosaics: the consequences of grazing vs. natural heterogeneity. *Ecol. Appl.*, 11, 1123–1140.
- Bestelmeyer, B. T. & Wiens, J. A. (2001). Local and regional-scale responses of ant diversity to a semiarid biome transition. *Ecography*, 24, 381–392.
- Biseau, J.-C., Quinet, Y., Deffernez, L. & Pasteels, J. M. (1997). Explosive food recruitment as a competitive strategy in the ant *Myrmica sabuleti* (Hymenoptera, Formicidae). *Insect. Soc.*, 44, 59–73.
- Bisevac, L. & Majer, J. D. (1999). Comparative study of ant communities of rehabilitated mineral sand mines and heathland, Western Australia. *Restor. Ecol.*, 7, 117–126.
- Bliss, P., Katzerke, A. & Moritz, R. F. A. (2000). *Formica exsecta* Nylander, 1846 (Hym., Formicidae) im Müritz-Nationalpark. *Ent. Nachr. Ber.*, 44, 283–284.
- Bliss, P., Schröder, H., Katzerke, A. & Moritz, R. F. A. (2001). Standort und Struktur eines Kolonieverbandes der Großen Kerbameise (*Formica exsecta*) im Müritz-Nationalpark (Hymenoptera, Formicidae). *Arch. Freunde Naturg. Mecklenb.*, 40, 5–23.
- Bliss, P. & Katzerke, A. (2002). *Formica exsecta* Nylander, 1846 in der Uckermark (Hym., Formicidae). *Ent. Nachr. Ber.*, 46, 189.
- Bliss, P. & Katzerke, A. (2004). Zur Bestandssituation und Gefährdung von *Formica foreli* im Müritz-Nationalpark nebst Anmerkungen zum Forschungsbedarf (Hym., Formicidae). *Ent. Nachr. Ber.*, 48, 19–22.
- Blüthgen, N., Verhaagh, M., Goitia, W., Jaffe, K., Morawetz, W. & Barthlott, W. (2000). How plants shape the ant community in the Amazonian rainforest canopy: the key role of extrafloral nectaries and homopteran honeydew. *Oecologia*, 125, 229–240.
- Blüthgen, N. & Fiedler, K. (2002). Interactions between weaver ants *Oecophylla smaragdina*, homopterans, trees and lianas in an Australian rain forest canopy. *J. Anim. Ecol.*, 71, 793–

- Blüthgen, N., Gottsberger, G. & Fiedler, K. (2004). Sugar and amino acid composition of ant-attended nectar and honeydew sources from an Australian rainforest. *Austral Ecol.*, 29, 418–429.
- Bokor, Z. (1993). Soil fauna studies in a beech forest II. Comparative studies on soil invertebrates in a forest, forest margin and a clear-cut area in Hungary. *Acta Biol. Szeged.*, 39, 77–91.
- Bonte, D., Dekoninck, W., Provoost, S., Cosijns, E. & Hoffmann, M. (2003). Microgeographical distribution of ants (Hymenoptera: Formicidae) in coastal dune grassland and their relation to the soil structure and vegetation. *Anim. Biol.*, 53, 367–377.
- Boomsma, J. J., Mabelis, A. A., Verbeek, M. G. M. & Los, E. C. (1987). Insular biogeography and distribution ecology of ants on the Frisian islands. *J. Biogeogr.*, 14, 21–37.
- Bönsel, A. & Busch, T. (2003). Beschreibung des bislang größten bekannten Vorkommens von *Formica (Coptoformica) foreli*. *Ameisenschutz aktuell*, 17, 74–83.
- Braman, S. K. & Pendley, A. F. (1993). Relative and seasonal abundance of beneficial arthropods in centipedegrass as influence by management practices. *J. Econ. Entomol.*, 86, 494–504.
- Braune, M. (1974). Zur Hymenopterenfauna von Agrozönosen und der Einfluß einer Herbizidbehandlung auf ihre Zusammensetzung. *Hercynia N.F., Leipzig*, 11, 299–323.
- Bretz, D. (1995). Ameisenschutz-Praxis (20). *Ameisenschutz aktuell*, 9, 60–65.
- Brian, M. V., Hibble, J., Kelly, A. F. (1966). The dispersion of ant species in a southern English heath. *J. Anim. Ecol.*, 35, 281–290.
- Brown, J. H., Davidson, D. W. (1977). Competition between seed-eating rodents and ants in

- desert ecosystems. *Science*, 196, 880–882.
- Brown, J. H., Davidson, D. W., Reichman, O. J. (1979). An experimental study of competition between seed-eating desert rodents and ants. *Amer. Zool.*, 19, 1129–1143.
- Bugrova, N. M., Reznikova, J. I. (1990). The state of *Formica polyctena* Foerst. (Hymenoptera, Formicidae) population in recreation forests. *Mem. zool.*, 44, 7–12.
- Burbidge, A. H., Leicester, K., McDavitt, S. & Majer, J. D. (1992). Ants as indicators of disturbance at Yanchep National Park, Western Australia. *J. R. Soc. West. Aust.*, 75, 89–95.
- Buschinger, A. (1975). Die Ameisenfauna des Bausenberges, der nordöstlichen Eifel und Voreifel (Hym., Formicidae) mit einer quantitativen Auswertung von Fallenfängen. *Beiträge Landespfllege Rhld.-Pfalz Beih.*, 4, 251–273.
- Cabrera, M., Jaffé, K., Goitia, W. & Osborn, F. (1998). Recovery of disturbed ecosystems as monitored by ant and vegetation diversity in forests and surrounding savannas of Venezuela. *Stud. Neotrop. Fauna & Environm.*, 33, 85–92.
- Cammell, M. E., Way, M. J. & Paiva, M. R. (1996). Diversity and structure of ant communities associated with oak, pine, eucalyptus and arable habitats in Portugal. *Insect. Soc.*, 43, 37–46.
- Castro, A. G., Queiroz, M. V. B. & Araújo, L. M. (1990). O papel do distúrbio na estrutura de comunidades de formigas (Hymenoptera, Formicidae). *Revta. bras. Ent.*, 34, 201–213.
- Cerdá, X. & Retana, J. (1988). Descripcion de la comunidad de hormigas de un prado sabanoide en Canet de Mar. *Ecología*, 2, 333–341.
- Cerdá, X. & Retana, J. (1990). Foraging strategies for insect corpses in an ant community (Hymenoptera: Formicidae). *Ses. Entom. ICHN-SCL*, 6, 109–118.
- Cerdá, X., Alsina, A., Retana, J. & Bosch, J. (1990). Coexistencia de tres especies simpatridas del genero *Camponotus* Mayr (Hymenoptera Formicidae). In Reyna Martinez, L. A.,

- Serrano, P. R. & Nevado, T. R. (Eds.), *Principios en etología* (pp. 117–130). Cordoba: Cajasur.
- Cerdá, X., Retana, J. & Cros, S. (1997). Thermal disruption of transitive hierarchies in Mediterranean ant communities. *J. Anim. Ecol.*, *66*, 363–374.
- Cerdá, X., Retana, J. & Manzaneda, A. (1998). The role of competition by dominants and temperature in the foraging of subordinate species in Mediterranean ant communities. *Oecologia*, *117*, 404–412.
- Cerdá, X., Retana, R. & Cros, S. (1998). Critical thermal limits in Mediterranean ant species: trade-off between mortality risk and foraging performance. *Funct. Ecol.*, *12*, 45–55.
- Cherix, D. & Gris, G. (1978). Relation et agressivité chez *Formica lugubris* Zett. dans le Jura. *Proc. 8th Meeting Soc. Ins. Sect. PES*, 7–14.
- Chew, A. E. & Chew, R. M. (1980). Body size as a determinant of small-scale distributions of ants in evergreen woodland Southeastern Arizona. *Insect. Soc.*, *27*, 189–202.
- Chew, R. M. (1987). Population dynamics of colonies of three species of ants in desertified grassland, Southeastern Arizona, 1958–1981. *Am. Midl. Nat.*, *118*, 177–188.
- Chew, R. M. (1995). Aspects of the ecology of three species of ants (*Myrmecocystus* spp., *Aphaenogaster* sp.) in desertified grassland in Southeastern Arizona, 1958–1993. *Am. Midl. Nat.*, *134*, 75–83.
- Clay, R. E. (1996). Ant fauna of a mangrove community in the Australian seasonal tropics, with particular reference to zonation. *Austr. J. Zool.*, *44*, 521–533.
- Cook, J. L. (2003). Conservation of biodiversity in an area impacted by the red imported fire ant, *Solenopsis invicta* (Hymenoptera: Formicidae). *Biodiv. Conserv.*, *12*, 187–195.
- Crist, T. O. & Wiens, J. A. (1996). The distribution of ant colonies in a semiarid landscape: implications for community and ecosystem processes. *Oikos*, *76*, 301–311.

- Cros, S., Cerdá, X. & Retana, J. (1997). Spatial and temporal variations in the activity patterns of Mediterranean ant communities. *Ecoscience*, 4, 269–278.
- Czechowski, W. (1976). Competition between *Formica exsecta* Nyl. and *Formica pressilabris* Nyl. (Hymenoptera, Formicidae). *Ann. Zool. Warsz.*, 33, 1–14.
- Czechowski, W. & Pisarski, B. (1988). Inter- and intraspecific competitive relations in *Camponotus ligniperdus* (Latr.) (Hymenoptera, Formicidae). *Ann. Zool. Warsz.*, 41, 355–381.
- Czechowski, W. & Pisarski, B. (1990). Ants (Hymenoptera, Formicidae) of the Vistula escarpment in Warsaw. *Fragm. faun.*, 33, 109–128.
- Czechowski, W., Czechowska, W. & Palmowska, J. (1990). Arboreal myrmecofauna of Warsaw parks. *Fragm. faun.*, 34, 37–45.
- Czechowski, W. & Pisarski, B. (1990). Ants (Hymenoptera, Formicoidea) of linden-oak-hornbeam forests and thermophilous oak forests of the Mazovian Lowland. 1. Nest density. *Fragm. faun.*, 34, 134–141.
- Czechowski, W., Pisarski, B. & Czechowski, W. (1990). Ants (Hymenoptera, Formicoidea) of moist meadows on the Mazovian Lowland. *Fragm. faun.*, 34, 47–60.
- Czechowski, W. (1991). Comparison of the myrmecofaunas (Hymenoptera, Formicoidea) of tree stands and lawns in Warsaw parks. *Fragm. faun.*, 35, 179–183.
- Czechowski, W., Pisarski, B. & Yamauchi, K. (1995). Succession of ant communities (Hymenoptera, Formicidae) in moist pine forests. *Fragm. faun.*, 38, 447–486.
- Czechowski, W. (1998). Succession of ant communities (Hymenoptera: Formicidae) in the moist pine forest habitat of Bialowieza Forest. *Parki Narodowe i Rezerwaty Przyrody*, 17, 55–62.
- Dauber, J. (1995). *Die Ameisen (Hymenopter: Formicidae) von Mainz*. Unpublished Master's

thesis. Mainz: Johannes Gutenberg-Universität Mainz.

Dauber, J. (2002). Nestabundanzen von Ameisen (Hymenoptera: Formicidae) in städtischen Kleinstrukturen. *Mainz. naturwiss. Arch., Beih.*, 22, 287–295.

Dauber, J. & Wolters, V. (2004). Edge-effects on ant community structure and species richness in an agricultural landscape. *Biodiv. Conserv.*, 13, 901–915.

Davidson, W. D. (1977). Species diversity and community organization in desert seed-eating ants. *Ecology*, 58, 711–724.

Davidson, D. W., Longino, J. T. & Snelling, R. R. (1988). Pruning of host plant neighbors by ants: an experimental approach. *Ecology*, 69, 801–808.

de Biseau, J.-C., Quinet, Y., Deffernez, L. & Pasteels, J. M. (1997). Explosive food recruitment as a competitive strategy in the ant *Myrmica sabuleti* (Hymenoptera, Formicidae). *Insect. Soc.*, 44, 59–73.

de Bruyn, G. J. & Mabelis, A. A. (1972). Predation and aggression as possible regulatory mechanisms in *Formica*. *Ekol. pol.*, 20, 93–101.

Dean, W. R. J., Milton, S. J. & Klotz, S. (1997). The role of ant nest-mounds in maintaining small-scale patchiness in dry grasslands in Central Germany. *Biodiv. Conserv.*, 6, 1293–1307.

Deffernez, L., Champagne, P., Verhaeghe, J.-C., Josens, G. & Loreau, M. (1990). Analysis of the spatio-temporal niche of foraging grassland ants in the field. *Insect. Soc.*, 37, 1–13.

Dejean, A., Masens, D., Kanika, K., Nsudi, M. & Gunumina, R. (1986). Les termites et les fourmis, animaux dominants de la faune du sol de plusieurs formations forestieres et herbeuses du Zaire. *Actes Coll. Insect. Soc.*, 3, 273–283.

Dejean, A. & Gibernau, M. (2000). A rainforest ant mosaic: the edge effect (Hymenoptera: Formicidae). *Sociobiology*, 35, 385–401.

- Dejean, A., McKey, D., Gibernau, M. & Belin, M. (2000). The arboreal ant mosaic in a Cameroonian rainforest (Hymenoptera: Formicidae). *Sociobiology*, 35, 403–423.
- Della Lucia, T. M .C., Loureiro, M. C., Chandler, L., Freire, J. A. H., Galvao, J. D. & Fernandes, B. (1982). Ordenacao de comunidades de Formicidae em quatro agroecossistemas em Vicoso, Minas Gerais. *Experientiae*, 28, 67–94.
- Deslippe, R. J. & Savolainen, R. (1994). Role of food supply in structuring a population of *Formica* ants. *J. Anim. Ecol.*, 63, 756–764.
- Dix, M. E. & Baxendale, F. (1997). Insect pests and arthropod predators associated with tree-turf landscapes. *J. Entomol. Sci.*, 32, 257–271.
- Doncaster, C. P. (1981). The spatial distribution of ants' nests on Ramsey Island, South Wales. *J. Anim. Ecol.*, 50, 195–218.
- Duelli, P. & Obrist, M. K. (1998). In search of the best correlates for local organismal biodiversity in cultivated areas. *Biodiv. Conserv.*, 7, 297–309.
- Dugas, D. P. (2001). The influence of arroyo edges on *Pogonomyrmex rugosus* nest distribution in the Chihuahuan Desert, New Mexico. *J. Arid Environ.*, 47, 153–159.
- Dyer, L. A. & Letourneau, D. K. (1999). Trophic cascades in a complex terrestrial community. *Proc. Natl. Acad. Sci. USA*, 96, 5072–5076.
- Eldridge, D. J. (1993). Effect of ants on sandy soils in semi-arid eastern Australia: Local distribution of nest entrances and their effect on infiltration of water. *Aust. J. Soil Res.*, 31, 509–518.
- Eldridge, D. J. & Pickard, J. (1994). Effects of ants on sandy soils in semi-arid eastern Australia: II. Relocation of nest entrances and consequences for bioturbation. *Aust. J. Soil Res.*, 32, 323–333.
- Elmes, G. W. (1974). The spatial distribution of a population of two ant species living in

- limestone grassland. *Pedobiologia*, 14, 412–418.
- Elmes, G. W. (1982). The phenology of five species of *Myrmica* (Hym. Formicidae) from South Dorset, England. *Insect. Soc.*, 29, 548–559.
- Elmes, G. W. & Wardlaw, J. C. (1982). Variations in populations of *Myrmica sabuleti* and *M. scabrinodis* (Formicidae: Hymenoptera) living in Southern England. *Pedobiologia*, 23, 90–97.
- Espadaler, X., Rey, S. & Bernal, V. (2004). Queen number in a supercolony of the invasive garden ant, *Lasius neglectus*. *Insect. Soc.*, 51, 232–238.
- Evans, E. W. (1991). Experimental manipulation of herbivores in native tallgrass prairie: Responses of aboveground arthropods. *Am. Midl. Nat.*, 125, 37–46.
- Feener, D. H., Schupp, E. W. (1998). Effect of treefall gaps on the patchiness and species richness of Neotropical ant assemblages. *Oecologia*, 116, 191–201.
- Feldhaar, H., Fiala, B., Hashim, R. & Maschwitz, U. (2003). Patterns of the *Crematogaster*-*Macaranga* association: The ant partner makes the difference. *Insect. Soc.*, 50, 9–19.
- Ferguson, S. H. (2001). Changes in trophic abundance of soil arthropods along a grass-shrub-forest gradient. *Can. J. Zool.*, 79, 457–464.
- Fernández-Escudero, I. & Tinaut, A. (1999). Factors determining nest distribution in the high-mountain ant *Proformica longiseta* (Hymenoptera Formicidae). *Ethol. Ecol. Evol.*, 11, 325–338.
- Fisher, B. L. (1997). A comparison of ant assemblages (Hymenoptera, Formicidae) on serpentine and non-serpentine soils in northern California. *Insect. Soc.*, 44, 23–33.
- Fisher, B. L. (1999). Ant diversity patterns along an elevational gradient in the Réserve Naturelle Intégrale d' Andohahela, Madagascar. *Fieldiana Zool.*, 94, 129–147.

- Floren, A. & Linsenmair, K. E. (1999). Changes in arboreal arthropod communities along a disturbance gradient. *Selbyana*, 20, 284–289.
- Floren, A. & Linsenmair, K. E. (2000). The influence of anthropogenic disturbances on the structure of arboreal arthropod communities. *Plant Ecol.*, 153, 153–167.
- Floren, A., Freking, A., Biehl, M. & Linsenmair, K. E. (2001). Anthropogenic disturbance changes the structure of arboreal tropical ant communities. *Ecography*, 24, 547–554.
- Floren, A., Biun, A. & Linsenmair, K. E. (2002). Arboreal ants as key predators in tropical lowland rainforest trees. *Oecologia*, 131, 137–144.
- Fonseca, C. R. & Ganade, G. (1996). Asymmetries, compartments and null interactions in an Amazonian ant-plant community. *J. Anim. Ecol.*, 65, 339–347.
- Fonseca, C. R. (1999). Amazonian ant-plant interactions and the nesting space limitation hypothesis. *J. Trop. Ecol.*, 15, 807–825.
- Foster, J. & Kettle, W. D. (1999). Distribution of a mound-building ant on native and restored prairies in Northeastern Kansas. *Prairie Nat.*, 31, 21–31.
- Fox, M. D. & Fox, B. J. (1982). Evidence for interspecific competition influencing ant species diversity in a regenerating heathland. In Buckley, R. C. (Ed.), *Ant-plant interactions in Australia* (pp. 99–110). The Hague: Dr. W. Junk Publishers.
- Fox, B. J. (1982). The influence of disturbance (fire, mining) on ant and small mammal species diversity in Australian heathland. *Proc. Symp. Dynamics and management of Mediterranean-type ecosystems*, 213–219.
- Fox, B. J., Fox, M. D. & Archer, E. (1985). Experimental confirmation of competition between two dominant species of *Iridomyrmex* (Hymenoptera: Formicidae). *Austr. J. Ecol.*, 10, 105–110.
- Francoeur, A. & Maldague, M. (1966). Classification des micromilieux de nidification des

- fourmis. *Naturaliste Can.*, 93, 473–478.
- Fründ, H.-C. (1990). Zur bodennahen Fauna in Straßen-Baumscheiben verschiedener Gestaltung. *Natur und Landschaft*, 65, 597–599.
- Gadagkar, R., Nair, P., Chandrashekara, K. & Bhat, D. M. (1993). Ant species richness and diversity in some selected localities in Western Ghats, India. *Hexapoda*, 5, 79–94.
- Gallé, L. (1967). Ecological and zoocoenological conditions of the Formicidae fauna at Tiszakürt. *Tiscia*, 3, 67–73.
- Gallé, L. (1969). Myrmecological investigations in the environs of Kisköre. *Tiscia*, 5, 87–95.
- Gallé, L. (1975) Factors stabilizing the ant populations (Hymenoptera: Formicidae) in the grass associations of the Tisza Basin. *Tiscia*, 10, 61–66.
- Gallé, L. (1978). Dispersion of the nests of an ant species (Hymenoptera: Formicidae). *Acta Biol. Szeged.*, 24, 105–109.
- Gallé, L. (1990). Assembly of sand-dune forest ant communities. *Mem. zool.*, 44, 1–6.
- Gallé, L. (1991). Structure and succession of ant assemblages in a north European sand dune area. *Holarct. Ecol.*, 14, 31–37.
- Gallé, L., Kovács, É. & Hevér, A. (1994). Pattern transformation of ant colonies in a successional sandy grassland. *Mem. zool.*, 48, 81–90.
- Gallé, L., Körmöczi, L. & Kerekes, J. (1998). Structure of ant assemblages in a Middle-European successional sand-dune area. *Tiscia*, 31, 19–28.
- Gallé, L. (1972). Formicidae populations of the ecosystems in the environs of Tiszafüred. *Tiscia*, 7, 59–68.
- Gallé, L. (1972). Study of ant-populations in various grassland ecosystems. *Acta Biol. Szeged*, 18, 159–164.

- Gallé, L. (1980). Dispersion of high density ant populations in sandy soil grassland ecosystems. *Acta Biol. Szeged*, 26, 129–135.
- Gallé, L. (1986). Habitat and niche analysis of grassland ants (Hymenoptera: Formicidae). *Entomol. Gener.*, 11, 197–211.
- Glaser, F. (1998). Die Ameisenfauna des Arzler Kalvarienberges bei Innsbruck (Tirol, Österreich). *Ber. nat.-med. Ver. Innsbruck*, 85, 257–286.
- Glaser, F., Kopf, T. & Steinberger, K.-H. (2003). Ameisen (Hymenoptera: Formicidae) im Frastanzer Ried und den Illauen (Vorarlberg, Österreich). Artenspektrum, Gefährdung und Schutzempfehlungen. *Vorarlberger Naturschau*, 13, 287–310.
- Golden, D. M. & Crist, T. O. (2000). Experimental effects of habitat fragmentation on rove beetles and ants: patch area or edge? *Oikos*, 90, 525–538.
- Golley, F. B. & Gentry, J. B. (1964). Bioenergetics of the Southern harvester ant, *Pogonomyrmex badius*. *Ecology*, 45, 217–225.
- Gomez, C., Casellas, D., Oliveras, J., Bas, J. M. (2003). Structure of ground-foraging ant assemblages in relation to land-use change in the northwestern Mediterranean region. *Biodiv. Conserv.*, 12, 2135–2146.
- Gordon, D. M. & Wagner, D. (1997). Neighborhood density and reproductive potential in harvester ants. *Oecologia*, 109, 556–560.
- Gotelli, N. J. (1996). Ant community structure: effects of predatory ant lions. *Ecology*, 77, 630–638
- Grey, E. J. (2002). Invertebrates of Mount McKay - a brief survey. *Vict. Nat.*, 119, 29–35.
- Gridina, T. I. (1990). Influence of *Formica polyctena* Foerst (Hymenoptera, Formicidae) on the distribution of predatory arthropods in forest ecosystems. *Mem. zool.*, 44, 13–20.

- Grossrieder, M. & Zettel, J. (1999). Auensukzession und Zonation im Rottensand (Pfynwald, Kt. VS). III. Wiederbesiedlung einer Überschwemmungsfläche durch Ameisen (Hymenoptera, Formicidae). *Mitt. Schweiz. Entomol. Ges.*, 72, 153–164.
- Haeseler, V. (1990). Zum Vorkommen von *Harpagoxenus sublaevis* (Nylander) in einem aufgeforsteten Hochmoorrest bei Oldenburg (Hymenoptera: Formicidae). *Braunschweig. naturkd. Schr.*, 3, 757–762.
- Hahn, D. A. & Tschinkel, W. R. (1997). Settlement and distribution of colony-founding queens of the arboreal ant, *Crematogaster ashmeadi*, in a longleaf pine forest. *Insect. Soc.*, 44, 323–336.
- Hanula, J. L. & New, K. C. P. (1996). A trap for capturing arthropods crawling up tree boles. *Southern Research Station Research Note SRS*, 3, 1–8.
- Harada, A. Y. & Adis, J. (1998). Ants obtained from trees of a "Jacareúba" (*Calophyllum brasiliense*) forest plantation in Central Amazonia by canopy fogging: first results. *Acta Amazon.*, 28, 309–318.
- Hawes, C., Stewart, A. J. A. & Evans, H. F. (2002). The impact of wood ants (*Formica rufa*) on the distribution and abundance of ground beetles (Coleoptera: Carabidae) in a Scots pine plantation. *Oecologia*, 131, 612–619.
- Herbers, J. M. (1985). Seasonal structuring of a north temperate ant community. *Insect. Soc.*, 32, 224–240.
- Herbers, J. M. (1989). Community structure in north temperate ants: temporal and spatial variation. *Oecologia*, 81, 201–211.
- Herbers, J. M. (1991). The population biology of *Tapinoma minutum* (Hymenoptera: Formicidae) in Australia. *Insect. Soc.*, 38, 195–204.
- Herbers, J. M. (1994). Structure of an Australian ant community with comparisons to North

- American counterparts (Hymenoptera: Formicidae). *Sociobiology*, 24, 293–306.
- Heterick, B. (1997). The interaction between the coastal brown ant, *Pheidole megacephala* (Fabricius), and other invertebrate fauna of Mt Coot-tha (Brisbane, Australia). *Aust. J. Ecol.*, 22, 218–221.
- Heterick, B. E., Casella, J., Majer, J. D. (2000). Influence of Argentine and coastal brown ant (Hymenoptera: Formicidae) invasions on ant communities in Perth gardens, Western Australia. *Urban Ecosyst.*, 4, 277–292.
- Hilzensauer, H. G. (1980). Zur Biologie und Ökologie von *Coptoformica exsecta* (Nylander, 1846) in der subalpinen Stufe des Patscherkofels (Tirol, Österreich). *Ber. nat.-med. Ver. Innsbruck*, 67, 173–184.
- Hinkley, S. & New, T. R. (1997). Pitfall trapping for surveying ant assemblages: lessons from a study at Mount Piper, Victoria. *Mem. Mus. Vict.*, 56, 369–376.
- Hoffmann, B. D., Andersen, A. N. & Hill, G. J. E. (1999). Impact of an introduced ant on native rain forest invertebrates: *Pheidole megacephala* in monsoonal Australia. *Oecologia*, 120, 595–604.
- Hoffmann, B. D., Griffiths, A. D. & Andersen, A. N. (2000). Responses of ant communities to dry sulfur deposition from mining emissions in semi-arid tropical Australia, with implications for the use of functional groups. *Austral Ecol.*, 25, 653–663.
- Hoffmann, B. D. (2000). Changes in ant species composition and community organisation along grazing gradients in semi-arid rangelands of the Northern Territory. *Rangel. J.*, 22, 171–189.
- Holway, D. A. (1998). Effect of Argentine ant invasions on ground-dwelling arthropods in northern California riparian woodlands. *Oecologia*, 116, 252–258.
- Holway, D. A. (1998). Factors governing rate of invasion: a natural experiment using

- Argentine ants. *Oecologia*, 115, 206–212.
- Holway, D. A. (1999). Competitive mechanisms underlying the displacement of native ants by the invasive Argentine ant. *Ecology*, 80, 238–251.
- Holway, D. A., Suarez, A. V. & Case, T. J. (2002). Role of abiotic factors in governing susceptibility to invasion: a test with Argentine ants. *Ecology*, 83, 1610–1619.
- Hosking, A. C. & Turner, V. (1997). Ants, disturbance and regeneration in eucalypt forests. *Mem. Mus. Vict.*, 56, 655–657.
- Höttinger, H., Schlick-Steiner, B. C. & Steiner, F. M. (2003). The alcon blue *Maculinea alcon* (Lepidoptera: Lycaenidae) in Eastern Austria: Status and conservation measures. *Ekologia (Bratislava)*, 22, 107–118.
- Hulugalle, N. R. (1995). Effects of ant hills on soil physical properties of a Vertisol. *Pedobiologia*, 39, 34–41.
- Human, K. G. & Gordon, D. M. (1997). Effects of Argentine ants on invertebrate biodiversity in Northern California. *Conserv. Biol.*, 11, 1242–1248.
- Igarashi, R. (1975). Faunal aspects in grassland. In Numata, M. (Ed.), *Ecological studies in Japanese grasslands with special reference to the IBP area - Productivity of terrestrial communities* (pp. 227–261). Tokyo: University of Tokyo Press.
- Jaffé, K. (1986). Control of *Atta* and *Acromyrmex* spp. in pine tree plantations in the Venezuelan Llanos. In Lofgren, C. S. & Meer, R. K. V. (Eds.), *Fire ants and leaf-cutting ants. Biology and management* (pp. 409–416). Boulder CO, London Westview Press.
- Jaffé, K. & Vilela, E. (1989). On nest densities of the leaf-cutting ant *Atta cephalotes* in tropical primary forest. *Biotropica*, 21, 234–236.
- Járdán, C., Gallé, L. & Margóczi, K. (1993). Ant assemblage composition in a successional hungarian sand dune area. *Tiscia*, 27, 9–15.

- Jennings, D. T., Houseweart, M. W. & Francoeur, A. (1986). Ants (Hymenoptera: Formicidae) associated with strip-clearcut and dense spruce-fir forests of Maine. *Can. Ent.*, 118, 43–50.
- Joger, H. G. (1988). Untersuchungen über die Tierwelt einer Stadtmauer. *Zool. Jb. Syst.*, 115, 69–91.
- Jusino-Atresino, R. & Phillips, S. A. (1994). Impact of red imported fire ants on the ant fauna of central Texas. In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 259–268). Boulder CO: Westview Press.
- Kajak, A. & Lukasiewicz, J. (1994). Do semi-natural patches enrich crop fields with predatory epigean arthropods. *Agr. Ecosyst. Environ.*, 49, 149–161.
- Kaneko, S. (2003). Impacts of two ants, *Lasius niger* and *Pristomyrmex pungens* (Hymenoptera: Formicidae), attending the brown citrus aphid, *Toxoptera citricidus* (Homoptera: Aphididae), on the parasitism of the aphid by the primary parasitoid, *Lysiphlebus japonicus* (Hymenoptera: Aphidiidae), and its larval survival. *Appl. Entomol. Zool.* 38, 347–357.
- Kaspari, M., O'Donnell, S. & Kercher, J. R. (2000). Energy, density, and constraints to species richness: ant assemblages along a productivity gradient. *Am. Nat.*, 155, 280–293.
- Kaspari, M., Pickering, J., Longino, J. T. & Windsor, D. (2001). The phenology of a Neotropical ant assemblage: evidence for continuous and overlapping reproduction. *Behav. Ecol. Sociobiol.*, 50, 382–390.
- Kaufmann, E., Malsch, A. K. F., Erle, M. & Maschwitz, U. (2003). Compound nesting of *Strumigenys* sp. (Myrmicinae) and *Diacamma* sp. (Ponerinae), and other nesting symbioses of myrmicine and ponerine ants in Southeast Asia. *Insect. Soc.*, 50, 88–97.
- Keals, N. & Majer, J. D. (1991). The conservation status of ant communities along Wubin-

- Perenjori Corridor. In Saunders, D. A. & Hobbs, R. J. (Eds.), *Nature Conservation 2: The role of corridors* (pp. 387–393). Chipping Norton, NSW: Surrey Beatty & Sons.
- Kennedy, T. A. (1998). Patterns of an invasion by argentine ants (*Linepithema humile*) in a riparian corridor and its effects on ant diversity. *Am. Midl. Nat.*, *140*, 343–350.
- Kerley, G. I. H. & Whitford, W. G. (2000). Impact of grazing and desertification in the Chihuahuan desert: plant communities, granivores and granivory. *Am. Midl. Nat.*, *144*, 78–91.
- King, J. R., Andersen, A. N. & Cutter, A. D. (1998). Ants as bioindicators of habitat disturbance: validation of the functional group model for Australia's humid tropics. *Biodiv. Conserv.*, *7*, 1627–1638.
- Klimes, L. & Sechterová, E. (1989). Epigaeic arthropods across an arable land and grassland interface. *Acta Entomol. Bohemoslov.*, *86*, 459–475.
- Klimetzek, D. (1981). Population studies on hill building wood-ants of the *Formica rufa*-Group. *Oecologia*, *48*, 418–421.
- Klimetzek, D. & Kobel-Lamparski, A. (1990). Die Ameisenfauna des Naturschutzgebietes "Isteiner Klotz". *Mitt. bad. Landesver. Naturkunde u. Naturschutz*, *15*, 145–158.
- Koponen, S. & Niemelä, P. (1995). Ground-living arthropods along pollution gradient in boreal pine forest. *Entomol. Fenn.*, *5*, 127–131.
- Koponen, S., Rinne, V. & Clayhills, T. (1997). Arthropods on oak branches in SW Finland, collected by a new trap type. *Entomol. Fenn.*, *3*, 177–183.
- Kotze, D. J. & Samways, M. J. (2001). No general edge effects for invertebrates at Afromontane forest/grassland ecotones. *Biodiv. Conserv.*, *10*, 443–466.
- Krushelnicky, P. D., Loope, L. L. & Joe, S. M. (2004). Limiting spread of a unicolonial invasive insect and characterization of seasonal patterns of range expansion. *Biol. Invas.*, *6*,

- Krzysztofiak, L. (1991). The effect of habitat pollution with heavy metals on ant populations and ant-hill soil. *Ekol. pol.*, 39, 181–202.
- Lehouck, V., Bonte, D., Dekoninck, W. & Maelfait, J.-P. (2004). The distribution of ant nests (Hymenoptera, Formicidae) in coastal grey dunes in Flanders (Belgium) and their relationship to myrmecochorous plants. *Belg. J. Zool.*, 134, 89–96.
- Lenoir, L. (2002). Can wood ants distinguish between good and bad food patches on the forest floor? *Eur. J. Soil Biol.*, 38, 97–102.
- Lenoir, L. (2003). Response of the foraging behaviour of red wood ants (*Formica rufa* group) to exclusion from trees. *Agric. Forest Entomol.*, 5, 183–189.
- Levings, S. C. (1983). Seasonal, annual, and among-site variation in the ground ant community of a deciduous tropical forest: some causes of patchy species distributions. *Ecol. Monogr.* 53, 435–455.
- Lobry de Bruyn, L. A. (1993). Ant composition and activity in naturally-vegetated and farmland environments on contrasting soils at Kellerberrin, Western Australia. *Soil Biol. Biochem.*, 25, 1043–1056.
- Lobry de Bruyn, L. A. & Conacher, A. J. (1994). The bioturbation activity of ants in agricultural and naturally vegetated habitats in semi-arid environments. *Aust. J. Soil. Res.* 32, 555–570.
- Lombarte i Carrera, A. (1986). *Contribucion al estudio de los formicidos (Hymenoptera, Formicidae) de la Serra de Collserola*. Unpublished master's thesis. Barcelona: Universidad Autónoma de Barcelona.
- Longcore, T. R. (1999). *Terrestrial arthropods as indicators of restoration success in coastal sage scrub*. Unpublished PhD thesis. Los Angeles: University of California.

- Lude, A., Reich, M. & Plachter, H. (1999). Life strategies of ants in unpredictable floodplain habitats of alpine rivers (Hymenoptera: Formicidae). *Entomol. Gener.*, 24, 75–91.
- Lynch, J. F., Balinsky, E. C., Vail, S. G. (1980). Foraging patterns in three sympatric forest ant species, *Prenolepis imparis*, *Paratrechina melanderi* and *Apaenogaster rufis* (Hymenoptera: Formicidae). *Ecol. Entomol.*, 5, 353–371.
- Lynch, J. F. (1981). Seasonal, successional, and vertical segregation in a Maryland ant community. *Oikos*, 37, 183–198.
- Mabelis, A. A. (1977). Artenreichtum von Ameisen in einigen Waldtypen. *Ber. Int. Symp. Int. Vereinig. Vegetationsk.*, 187–208.
- Mabelis, A. A. (1979). Wood ant wars. *Neth. J. Zool.*, 29, 451–620.
- Mabelis, A. A. (1979). Nest splitting by the red wood ant (*Formica polyctena* Foerster). *Neth. J. Zool.*, 29, 109–125.
- Mabelis, A. A. (1980). Distribution of red wood ants (*Formica polyctena* Först.) over the foraging area of their nest, and the influence of a conspecific neighbouring population. *Meijendel-comité*, 48, 221–232.
- Mabelis, A. A. (1994). Flying as a survival strategy for wood ants in a fragmented landscape (Hymenoptera, Formicidae). *Mem. Zool.*, 48, 147–170.
- Mabelis, A. A. & Korczynska, J. (2001). Dispersal for survival: Some observations on the trunk ant (*Formica truncorum* Fabricius). *Neth. J. Zool.*, 51, 299–321.
- Madden, K. E. & Fox, B. J. (1997). Arthropods as indicators of the effects of fluoride pollution on the succession following sand mining. *J. Appl. Ecol.*, 34, 1239–1256.
- Maggini, R., Guisan, A. & Cherix, D. (2002). A stratified approach for modeling the distribution of a threatened ant species in the Swiss National Park. *Biodiv. Conserv.*, 11, 2117–2141.

- Majer, J. D. (1972). The ant mosaic in Ghana cocoa farms. *Bull. Entomol. Res.*, 62, 151–160.
- Majer, J. D. (1983). Ants: bio-indicators of minesite rehabilitation, land use, and land conservation. *Environ. Manag.*, 7, 375–383.
- Majer, J. D. (1984). Recolonisation by ants in rehabilitated open-cut mines in Northern Australia. *Reclam. Reveg. Res.*, 2, 279–298.
- Majer, J. D. (1985). Recolonization by ants of rehabilitated mineral sand mines on North Stradbroke Island, with particular reference to seed removal. *Aust. J. Ecol.*, 10, 31–48.
- Majer, J. D. (1993). Comparison of the arboreal ant mosaic in Ghana, Brazil, Papua New Guinea and Australia - its structure and influence on arthropod diversity. In LaSalle, J. & Gauld, I. D. (Eds.), *Hymenoptera and biodiversity* (pp. 115–141). Wallingford: CAB International.
- Majer, J. D., Camer-Pesci, P. (1991). Ant species in tropical Australian tree crops and native ecosystems – is there a mosaic? *Biotropica*, 23, 173–181.
- Majer, J. D. & Kock, A. E. (1992). Ant recolonization of sand mines near Richards Bay, South Africa: an evaluation of progress with rehabilitation. *Suid-Afrik. Tydskr. Wetenskap.*, 88, 31–36.
- Majer, J. D., Recher, H. F. & Postle, A. C. (1994). Comparison of arthropod species richness in Eastern and Western Australian canopies: A contribution to the species number debate. *Mem. Queensl. Mus.*, 36, 121–131.
- Majer, J. D. (1994). Spread of Argentine ants (*Linepithema humile*), with special reference to Western Australia. In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 163–173). Boulder CO: Westview Press.
- Majer, J. D. & Nichols, O. G. (1998). Long-term recolonization patterns of ants in Western Australian rehabilitated bauxite mines with reference to their use as indicators of restoration

success. *J. Appl. Ecol.*, 35, 161–182.

Majer, J. D., Recher, H. F., Heterick, B. E. & Postle, A. C. (2002). The canopy, bark, soil and litter invertebrate fauna of the Darling Plateau and adjacent woodland near Perth, Western Australia, with reference to the diversity of forest and woodland invertebrates. *Pac. Conserv. Biol.*, 7, 229–239.

Major, R. E., Smith, D., Cassis, G., Gray, M. & Colgan, D. J. (1999). Are roadside strips important reservoirs of invertebrate diversity? A comparison of the ant and beetle faunas of roadside strips and large remnant woodlands. *Aust. J. Zool.*, 47, 611–624.

Malicky, H. (1968). Der Einfluss andauernder Beweidung auf die Kleintierfauna der Hutweiden im Seewinkel (Burgenland): Allgemeines und Formicidae. *Wiss. Arbeiten Bgld.*, 40, 58–64.

Malicky, H. (1968). Über den jahreszeitlichen Aktivitätsverlauf der Ameisen (Formicidae, Hymenoptera) auf den Apetloner Hutweiden (Burgenland). *Wiss. Arbeiten Bgld.*, 40, 79–82.

Markó, B. & Czechowski, W. (2004). *Lasius psammophilus* Seifert and *Formica cinerea* Mayr (Hymenoptera: Formicidae) on sand dunes: conflicts and coexistence. *Ann. Zool. Warsz.*, 54, 365–378.

Mattoni, R., Longcore, T. & Novotny, V. (2000). Environmental auditing. Arthropod monitoring for fine-scale habitat analysis: a case study of the El Segundo sand dunes. *Environ. Manag.*, 25, 445–452.

McGlynn, T. P. & Kirksey, S. E. (2000). The effects of food presentation and microhabitat upon resource monopoly in a ground-foraging ant (Hymenoptera: Formicidae) community. *Rev. Biol. Trop.*, 48, 629–642.

McIntyre, N. E., Rango, J., Fagan, W. F. & Faeth, S. H. (2001). Ground arthropod community structure in a heterogeneous urban environment. *Landscape Urban Plan.*, 52, 257–274.

Meier, R. E. (1994). Coexisting patterns and foraging behavior of introduced and native ants (Hymenoptera Formicidae) in the Galapagos Islands (Ecuador). In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 44–62). Boulder CO: Westview Press.

Meyer, H. & Reinke, H.-D. (1996). Veränderungen in der biozönotischen Struktur der Wirbellosenfauna von Salzwiesen durch unterschiedliche Beweidungsintensitäten mit Schafen. *Faun.-Ökol. Mitt.*, 7, 109–151.

Mody, K. & Linsenmair, K. E. (2003). Finding its place in a competitive ant community: leaf fidelity of *Camponotus sericeus*. *Insect. Soc.*, 50, 191–198.

Morris, T. I., Campos, M., Kidd, N. A. C., Jervis, M. A. & Symondson, W. O. C. (1999). Dynamics of the predatory arthropod community in Spanish olive groves. *Agric. Forest Entomol.*, 1, 219–228.

Morton, S. R. (1988). Comparative structure of harvester ant communities in arid Australia and North America. *Ecol. Monogr.*, 58, 19–38.

Moya-Ragoza, G. & Nault, L. R. (2000). Obligatory mutualism between *Dalbulus quinquenotatus* (Homoptera: Cicadellidae) and attendant ants. *Ann. Entomol. Soc. Am.*, 93, 929–940.

Münch, W. (1983). Die Ameisen der Tübinger Neuhalde, eine faunistisch-ökologische Bestandsaufnahme der Nestdichten. *Veröff. Naturschutz Landschaftspflege Bad.-Württ.*, 57/58, 305–324.

Münch, W. (1995). Vorkommen und Verbreitung der Ameisen im Gebiet des "Schmiechener Sees". *Beih. Veröff. Naturschutz Landschaftspflege Bad.-Württ.*, 78, 435–475.

Nash, M. S., Whitford, W. G., van Zee, J. & Havstad, K. (1998). Monitoring changes in stressed ecosystems using spatial patterns of ant communities. *Environ. Monit. Assess.*, 51,

201–210.

Nash, M. S., Whitford, W. G., van Zee, J. & Havstad, K. M. (2000). Ant (Hymenoptera: Formicidae) responses to environmental stressors in the Northern Chihuahuan Desert.

Environ. Entomol., 29, 200–206.

Nash, M. S., Whitford, W. G., Bradford, D. F., Franson, S. E., Neale, A. C. & Heggem, D. T. (2001). Ant communities and livestock grazing in the Great Basin, U.S.A. *J. Arid Environ.*, 49, 695–710.

Nash, M. S. & Whitford, W. G. (2001). Ants as biological indicators for monitoring changes in arid environments: lessons for monitoring protected areas. Research and management options for protected areas. *Proc. First Int. Symp. Workshop Arid Zone Environ.*, 105–121.

Negi, H. R. & Gadgil, M. (2002). Cross-taxon surrogacy of biodiversity in the Indian Garhwal Himalaya. *Biol. Conserv.*, 105, 143–155.

New, K. C. & Hanula, J. L. (1998). Effect of time elapsed after prescribed burning in longleaf pine stands on potential prey of the red-cockaded woodpecker. *South. J. Appl. For.*, 22, 175–183.

Nielsen, M. G. (1974). Number and biomass of worker ants in a sandy heath area in Denmark. *Nat. Jutl.*, 17, 93–95.

Nielsen, M. G., Skyberg, N. & Winther, L. (1976). Studies on *Lasius flavus* F. (Hymenoptera, Formicidae): I. Population density, biomass, and distribution of nests. *Ent. Meddr.*, 44, 65–75.

Nielsen, M. G. (1981). The ant fauna on the high salt marsh. In Smit, C. J., Hollander, J., Wingerden, W. K. R. E. & Wolff, W. J. (Eds.), *Terrestrial and freshwater fauna of the Wadden Sea area* (pp. 68–70). Leiden: Stichting Veth tot Steun aan Waddenonderzoek.

Nielsen, M. G. (1986). Ant nests on tidal meadows in Denmark. *Entomol. Gener.*, 11, 191–

- O'Dowd, D. J., Green, P. T. & Lake, P. S. (2003). Invasional "meltdown" on an oceanic island. *Ecol. Lett.*, 6, 812–817.
- Obermann, H.-W. (2000). *Die Trockenmauer - der verborgene Biotop. Besiedlung des Ökosystems durch ausgewählte Tiergruppen, dargestellt am Beispiel zweier Weinbergslagen an der Mosel*. Bad Dürkheim: Eigenverlag der Pollichia.
- Ofer, J. (1976). Societies of ant species in Israel. *Israel J. Zool.*, 25, 217–218.
- Oliver, I., Nally, R. M. & York, A. (2000). Identifying performance indicators of the effects of forest management on ground-active arthropod biodiversity using hierarchical partitioning and partial canonical correspondence analysis. *Forest Ecol. Manag.*, 139, 21–40.
- Orr, M. R., Dahlsten, D. L. & Benson, W. W. (2003). Ecological interactions among ants in the genus *Linepithema*, their phorid parasitoids, ant ant competitors. *Ecol. Entomol.*, 28, 203–210.
- Osborn, F., Goitia, W., Cabrera, M. & Jaffé, K. (1999). Ants, plants and butterflies as diversity indicators: comparisons between strata at six forest sites in Venezuela. *Stud. Neotrop. Fauna & Environm.*, 34, 59–64.
- Paarmann, W. & Kerck, K. (1997). Advances in using the canopy fogging technique to collect living arthropods from tree-crowns. In Stork, N. E., Adis, J. & Didham, R. K. (Eds.), *Canopy arthropods* (pp. 53–66). London: Chapman & Hall.
- Peck, S. L., McQuaid, B. & Campbell, C. L. (1998). Using ant species (Hymenoptera: Formicidae) as a biological indicator of agroecosystem condition. *Environ. Entomol.*, 27, 1102–1110.
- Pedersen, J. S. & Boomsma, J. J. (1999). Genetic analysis of colony structure in polydomous

- and polygynous ant populations. *Biol. J. Linn. Soc.*, 66, 115–144.
- Pellens, R. & Garay, I. (2000). Edaphic macroarthropod communities in fast-growing plantations of *Eucalyptus grandis* Hill ex Maid (Myraceae) and *Acacia mangium* Wild (Leguminosae) in Brazil. *Eur. J. Soil Biol.*, 35, 77–89.
- Peng, R. K., Christian, K. & Gibb, K. (1998). The effect of non-crop vegetation on the insect pests and their natural enemies in cashew (*Anacardium occidentale* L.) plantations. *Plant Prot. Quart.*, 13, 16–20.
- Perfecto, I. & Snelling, R. (1995). Biodiversity and the transformation of a tropical agroecosystem: ants in coffee plantations. *Ecol. Appl.*, 5, 1084–1097.
- Perfecto, I. & Vandermeer, J. (1996). Microclimatic changes and the indirect loss of ant diversity in a tropical agroecosystem. *Oecologia*, 108, 577–582.
- Perfecto, I., Vandermeer, J., Hanson, P. & Cartín, V. (1997). Arthropod biodiversity loss and the transformation of a tropical agro-ecosystem. *Biodiv. Conserv.*, 6, 935–945.
- Petal, J., Jakubczyk, H. & Wojcik, Z. (1970). Influence des fourmis sur les modifications des sols et des plantes dans les milieux de prairie. In Phillipsen, J. (Ed.), *Methods of study in soil ecology* (pp. 235–239). New York: UNIPUB.
- Petal, J. M. (1976). The effect of mineral fertilization on ant populations in meadows. *Pol. Ecol. Stud.*, 2, 209–218.
- Petal, J. (1980). Ant populations, their regulation and effect on soil in meadows. *Ekol. pol.*, 28, 297–326.
- Petal, J. (1980). The effect of industrial pollution of Silesia on populations of ants. *Pol. Ecol. Stud.*, 6, 665–672.
- Petal, J. (1981). Intraspecific competition as a way of adaptation to food resources in an ant population. *Ekol. pol.*, 29, 421–430.

Petal, J. (1991). Diversity of ant communities in peatbogs of various origin and period under management. *Pol. Ecol. Stud.*, 17, 249–265.

Petal, J., Chmielewski, K., Czepinska-Kaminska, D., Konecka-Betley, K. & Kulinska, D. (1992). Ant communities in relation to changes in some properties of hydrogenic soils differentially transformed. *Ekol. pol.*, 40, 553–576.

Petrov, I. Z. & Gallé, L. (1986). Nest distribution of the ant *Cataglyphis c. aenescens* (Nyl.) (Hymenoptera, Formicidae) in Deliblato sandy area (Yugoslavia). *Ekologija*, 21, 135–148.

Peveling, R., Attignon, S., Langewald, J. & Ouambama, Z. (1999). An assessment of the impact of biological and chemical grasshopper control agents on ground-dwelling arthropods in Niger, based on presence/absence sampling. *Crop. Prot.*, 18, 323–339.

Peveling, R. (2000). Impact of locust control on non-target arthropods - environmental monitoring, risk assessment and risk management. *Mitt. Dtsch. Ges. Allg. Angew. Ent.*, 12, 149–154.

Peveling, R., Weyrich, J. & Müller, P. (1994). Side-effects of botanicals, insect growth regulators and entomopathogenic fungi on epigeal non-target arthropods in locust control. In Krall, S. & Wilps, H. (Eds.), *New trends in locust control* (pp. 147–176). Rossdorf: TZ-Verlag.

Pfeiffer, M., Chimedregzen, L. & Ulykpan, K. (2003). Community organization and species richness of ants (Hymenoptera/Formicidae) in Mongolia along an ecological gradient from steppe to Gobi desert. *J. Biogeogr.*, 30, 1921–1935.

Phillips, S. A., Rogers, W. M., Wester, D. B. & Chandler, L. (1987). Ordination analysis of ant faunae along the range expansion front of the red imported fire ant in South-central Texas. *Texas J. Agric. Nat. Res.*, 1, 11–15.

Pik, A. J., Dangerfield, J. M., Bramble, R. A., Angus, C. & Nipperess, D. A. (2002). The use

- of invertebrates to detect small-scale habitat heterogeneity and its application to restoration practices. *Environ. Monit. Assess.*, 75, 179–199.
- Pisarski, B. (1971). Zoologische Charakteristik der Landbiotopen vom Bieszczady-Gebirge. *Fragm. Faun.*, 17, 23–30.
- Pisarski, B. & Czechowski, W. (1978). Influence de la pression urbaine sur la myrmécofaune. *Mem. Zool.*, 29, 109–128.
- Pisarski, B. (1982). La pression des populations de *Formica (Coptoformica) exsecta* Nyl. sur le milieu. *Mem. Zool.*, 38, 261–271.
- Pisarski, B. & Vepsäläinen, K. (1989). Competition hierarchies in ant communities (Hymenoptera, Formicidae). *Ann. Zool. Warsz.*, 42, 321–329.
- Pontin, A. J. (1961). Population stabilization and competition between the ants *Lasius flavus* (F.) and *L. niger* (L.). *J. Anim. Ecol.*, 30, 47–54.
- Pontin, A. J. (1978). The numbers and distribution of subterranean aphids and their exploitation by the ant *Lasius flavus* (Fabr.). *Ecol. Entomol.*, 3, 203–207.
- Pontin, A. J. (1997). Intraspecific nest spacing and the diversity of species in an ant community. *Entomol. Mon. Mag.*, 133, 1–8.
- Porter, S. D. & Savignano, D. A. (1990). Invasion of polygyne fire ants decimates native ants and disrupts arthropod community. *Ecology*, 71, 2095–2106.
- Puntila, P., Haila, Y., Pajunen, T. & Tukia, H. (1991). Colonisation of clearcut forests by ants in the southern Finnish taiga: a quantitative survey. *Oikos*, 61, 250–262.
- Puntila, P., Koponen, S. & Saaristo, M. (1994). Colonisation of a burned mountain-birch forest by ants (Hymenoptera, Formicidae) in subarctic Finland. *Mem. Zool.*, 48, 193–206.
- Puntila, P., Haila, Y., Niemelä, J. & Pajunen, T. (1994). Ant communities in fragments of

- old-growth taiga and managed surroundings. *Ann. Zool. Fennici.*, 31, 131–144.
- Punttila, P. (1996). Succession, forest fragmentation, and the distribution of wood ants. *Oikos*, 75, 291–298.
- Punttila, P., Haila, Y. & Tukia, H. (1996). Ant communities in taiga clearcuts: habitat effects and species interactions. *Ecography*, 19, 16–28.
- Punttila, P. & Haila, Y. (1996). Colonisation of a Burned Forest by Ants in the Southern Finnish Boreal Forest. *Silva Fenn.*, 30, 421–435.
- Quinn, M. A., Kepner, R. L., Walgenbach, D. D., Foster, R. N., Bohls, R. A., Pooler, P. D., Reuter, K. C. & Swain, J. L. (1993). Grasshopper Stages of Development as Indicators of Nontarget Arthropod Activity: Implications for Grasshopper Management Programs on Mixed-Grass Rangeland. *Environ. Entomol.*, 22, 532–540.
- Rabeling, C., Verhaagh, M. & Engels, W. (2003). Nestdichte nicht blattschneidender, pilzzüchtender Ameisen (Formicidae, Myrmicinae, Attini) im amazonischen terra firme Regenwald und auf einer benachbarten Agroforstfläche. *Proc. Meeting German speaking IUSSI section, Regensburg 2003*, 36–37.
- Read, J. L. & Andersen, A. N. (2000). The value of ants as early warning bioindicators: responses to pulsed cattle grazing at an Australian arid zone locality. *J. Arid Environ.*, 45, 231–251.
- Richter, K. & Klausnitzer, B. (1986). Zur Ameisenfauna unterschiedlich urban beeinflußter Ruderalstellen im Gebiet von Leipzig (Hym., Formicidae). *Entomol. Nachr. Ber.*, 30, 115–120.
- Roberts, D. L., Cooper, R. J. & Petit, L. J. (2000). Use of premontane moist forest and shade coffee agroecosystems by army ants in Western Panama. *Conserv. Biol.*, 14, 192–199.
- Rohe, W. (1992). Vergleichende Untersuchungen zur Ameisenfauna (Hymenoptera:

Formicidae) von Streuobstwiesen im Nordpfälzer Bergland. *Beitr. Landespfl. Rheinland-Pfalz*, 15, 495–529.

Rojas, P. & Fragoso, C. (2000). Composition, diversity, and distribution of a Chihuahuan desert ant community (Mapimí, México). *J. Arid Environ.*, 44, 213–227.

Rosengren, R., Vepsäläinen, K. & Wuorenrinne, H. (1979). Distribution, nest densities, and ecological significance of wood ants (the *Formica rufa* group) in Finland. *Bull. SROP*, 11, 183–213.

Rosengren, R., Cherix, D. & Pamilo, P. (1986). Insular ecology of the red wood ant *Formica truncorum* Fabr. II. Distribution, reproductive strategy and competition. *Mitt. Schweiz. Entomol. Ges.*, 59, 63–94.

Rosengren, R. (1986). Competition and coexistence in an insular ant community - a manipulation experiment (Hymenoptera: Formicidae). *Ann. Zool. Fennici*, 23, 297–302.

Roth, D. S. & Perfecto, I. (1994). The effects of management systems on ground-foraging ant diversity in Costa Rica. *Ecol. Appl.*, 41, 423–436.

Samways, M. J. (1983). Community structure of ants (Hymenoptera: Formicidae) in a series of habitats associated with citrus. *J. Appl. Ecol.*, 20, 833–847.

Samways, M. J. (1990). Ant assemblage structure and ecological management in citrus and subtropical fruit orchards in Southern Africa. In Vander Meer, R. K., Jaffé, K. & Cedeno, A. (Eds.), *Applied myrmecology* (pp. 570–587). Boulder, San Francisco & Oxford: Westview Press.

Samways, M. J., Osborn, R. & Carliel, F. (1997). Effect of a highway on ant (Hymenoptera: Formicidae) species composition and abundance, with a recommendation for roadside verge width. *Biodiv. Conserv.*, 6, 903–913.

Sanders, N. J. & Gordon, D. M. (2000). The effects of interspecific interactions on resource

- use and behavior in a desert ant. *Oecologia*, 125, 436–443.
- Sanders, N. J. & Gordon, D. M. (2003). Resource-dependent interactions and the organization of desert ant communities. *Ecology*, 84, 1024–1031.
- Savolainen, R. & Vepsäläinen, K. (1988). A competition hierarchy among boreal ants: impact on resource partitioning and community structure. *Oikos*, 51, 135–155.
- Savolainen, R. & Vepsäläinen, K. (1989). Niche differentiation of ant species within territories of the wood ant *Formica polyctena*. *Oikos*, 56, 3–16.
- Savolainen, R., Vepsäläinen, K. & Wuorenrinne, H. (1989). Ant assemblages in the taiga biome: testing the role of territorial wood ants. *Oecologia*, 81, 481–486.
- Schatz, B. & Hossaert-McKey, M. (2003). Interactions of the ant *Crematogaster scutellaris* with the fig/fig wasp mutualism. *Ecol. Entomol.*, 28, 359–368.
- Schlaghamerský, J. (1999). The impact of flooding on saproxylic beetles and ants in South Moravian floodplain forests. In: Tajovsky, K. & Pizl, V. (Eds.), *Soil zoology in Central Europe* (pp. 277–287). Ceske Budejovice: Institute of Soil Biology, Academy of Sciences of the Czech Republic.
- Schlaghamerský, J. (1999). The impact of flooding on saproxylic beetles and ants in South Moravian floodplain forests. *Soil Zoology in Central Europe. Proc. 5th Eur. Workshop on Soil Zoology*, 277–288.
- Schlick-Steiner, B. C., Steiner, F. M. & Höttlinger, H. (2002). Gefährdung und Schutz des Kreuzenzian-Ameisen-Bläulings *Maculinea rebeli* in Niederösterreich und Burgenland (Lepidoptera, Lycaenidae). *Linzer biol. Beitr.*, 34, 349–376.
- Schlick-Steiner, B. C., Steiner, F. M. & Moder, K. (2003). Ant nests in linear habitats: A new technique for estimating the density of sedentary organisms. Simulation and case study. *Sociobiology*, 42, 163–174.

- Schooley, R. L., Bestelmeyer, B. T. & Kelly, J. F. (2000). Influence of small-scale disturbances by kangaroo rats on Chihuahuan desert ants. *Oecologia*, 125, 142–149.
- Schooley, R. L. & Wiens, J. A. (2003). Spatial patterns, density dependence, and demography in the harvester ant, *Pogonomyrmex rugosus*, in semi-arid grasslands. *J. Arid Environ.*, 53, 183–196.
- Seifert, B. (1982). Die Ameisenfauna (Hymenoptera, Formicidae) einer Rasen-Wald-Catena im Leutratal bei Jena. *Abh. Ber. Naturkundemus. Görlitz*, 56, 1–18.
- Seifert, B. (1986). Vergleichende Untersuchungen zur Habitatwahl von Ameisen (Hymenoptera: Formicidae) im mittleren und südlichen Teil der DDR. *Abh. Ber. Naturkundemus. Görlitz*, 59, 1–124.
- Serrano, J. M., Acosta, F. J., Zorrilla, J. M. & Alvarez, M. (1988). El tiempo como variable relativa durante la sucesion de mirmecocenosis en eriales. *Stud. ecol.*, 5, 271–285.
- Serrano, J. M., Acosta, F. J. & López, F. (1993). Belowground space occupation and partitioning in an ant community during succession. *Eur. J. Entomol.*, 90, 149–158.
- Siepel, H., Meijer, J., Mabelis, A. A. & Boer, M. H. (1989). A tool to assess the influence of management practices on grassland surface macrofaunas. *J. Appl. Ent.*, 108, 271–290.
- Soares, S. M. & Schoereder, J. H. (2001). Ant-nest distribution in a remnant of tropical rainforest in southeastern Brazil. *Insect. Soc.*, 48, 280–286.
- Sonnenburg, H. & Behr, D. (1995). Die Ameisenfauna (Hymenoptera, Formicidae) der Streuobstwiesen bei Wehlen (Kreis Bernkastel-Wittlich) mit weiteren Funden aus dem Moselgebiet. *Dendrocopos*, 22, 90–105.
- Sonnenburg, H. (1996). Die Ameisenfauna (Hymenoptera, Formicidae) von Streuobstwiesen bei Rastatt (Baden). *Carolinea*, 54, 139–148.
- Söntgen, M. (1989). Ameisen (Formicidae). In Schulte, W., Fründ, H. C., Graefe, U.,

Ruszkowski, B., Söntgen, M., Voggenreiter, V. & Weritz, N. (Eds.), *Zur Biologie städtischer Böden – Beispielraum Bonn-Bad Godesberg* (pp. 118–119). Greven: Kilda-Verlag.

Stankiewicz, A. & Sieleznew, M. (2002). Host specificity of *Maculinea teleius* Bgstr. and *M. nausithous* Bgstr. (Lepidoptera: Lycaenidae). The new insight. *Ann. Zool. Warsz.*, 52, 403–408.

Steiner, F. M., Schlick-Steiner, B. C., Schödl, S., Espadaler, X., Seifert, B., Christian, E. & Stauffer, C. (2004). Phylogeny and bionomics of *Lasius austriacus* (Hymenoptera, Formicidae). *Insect. Soc.*, 51, 24–29.

Suarez, A. V., Bolger, D. T. & Case, T. J. (1998). Effects of fragmentation and invasion on native ant communities in coastal southern California. *Ecology*, 79, 2041–2056.

Suzzoni, J. P., Kenne, M. & Dejean, A. (1994). The ecology and distribution of *Myrmicaria opaciventris*. In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 133–150). Boulder CO: Westview Press.

Tartally, A. (2000). Notes on the coexistence of the supercolonial *Lasius neglectus* van Loon, Boomsma et Andrásfalvy 1990 (Hymenoptera: Formicidae) with other ant species. *Tiscia*, 32, 43–46.

Tassone, R. A. & Majer, J. D. (1997). Abundance of arthropods in tree canopies of *Banksia* woodland on the Swan Coastal Plain. *J. Rl. Soc. West Australia*, 80, 281–286.

Tennant, L. E. (1994). The ecology of *Wasmannia auropunctata* in primary tropical rainforest in Costa Rica and Panama. In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 80–90). Boulder CO: Westview Press.

Terayama, M. & Murata, K. (1990). Effects of area and fragmentation of forests for nature conservation: Analysis by ant communities. *Bull. Biogeog. Soc. Japan*, 45, 11–17.

- Terayama, M. (1993). Structure of ant communities in East Asia. II. Species and nest densities. *Bull. Biogeogr. Soc. Japan*, 48, 51–57.
- Theobald-Ley, S. & Horstmann, K. (1990). Die Ameisenfauna (Hymenoptera, Formicidae) von Windwurfflächen und angrenzenden Waldhabitaten im Nationalpark Bayerischer Wald. *Waldhygiene*, 18, 93–118.
- Thomas, M. L., Parry, L. J., Allan, R. A. & Elgar, M. A. (1999). Geographic affinity, cuticular hydrocarbons and colony recognition in the Australian meat ant *Iridomyrmex purpureus*. *Naturwissenschaften*, 86, 87–92.
- Thomas, J. A. (2002). Larval niche selection and evening exposure enhance adoption of a predacious social parasite, *Maculinea arion* (large blue butterfly), by *Myrmica* ants. *Oecologia*, 132, 531–537.
- Tobin, J. E. (1997). Competition and coexistence of ants in a small patch of rainforest canopy in Peruvian Amazonia. *J. New York Entomol. Soc.*, 105, 105–112.
- Touyama, Y., Ogata, K. & Sugiyama, T. (2003). The Argentine ant, *Linepithema humile*, in Japan: Assessment of impact on species diversity of ant communities in urban environments. *Entomol. Sci.*, 6, 57–62.
- Travan, J. (1984). Bestandsaufnahme der Waldameisennester in den Staatsforsten Unterfrankens, unter Berücksichtigung ökologischer Gesichtspunkte. *Waldhygiene*, 15, 65–94.
- Travan, J. (1990). Bestandsaufnahme der Waldameisennester im Staatswald Oberbayerns Teil I: Flachlandforstämter. *Waldhygiene*, 18, 119–142.
- Travan, J. (1996). Die Bestandsaufnahme von Waldameisenvölkern im bayerischen Flach- und Hügelland als Grundlage für deren Schutz durch Forstleute und freiwillige Helfer. *Ameisenschutz aktuell*, 10, 25–32.

Travan, J. (1997). Erfahrungen mit großflächigen Bestandsaufnahmen von Waldameisennestern in Bayern. *Ameisenschutz aktuell*, 11, 11–25.

Travan, J. (1997). Bestandsaufnahme der Waldameisenester im Staatswald Oberbayerns Teil II: Hochgebirgsforstämter. *Ameisenschutz aktuell*, 11, 69–108.

Travan, J. (1998). Über den Einfluß von Standortfaktoren auf die Besiedlung des bayrischen Hochgebirges durch Waldameisen (*Formica* spp.) (Hymen., Formicidae). *Anz. Schädlingskde., Pflanzenschutz, Umweltschutz*, 71, 105–109.

Tschinkel, W. R. & Hess, C. A. (1999). Arboreal ant community of a pine forest in Northern Florida. *Ann. Entomol. Soc. Am.*, 92, 63–70.

Tshiguvho, T. E., Dean, W. R. J. & Robertson, H. G. (1999). Conservation value of road verges in the semi-arid Karoo, South Africa: ants (Hymenoptera: Formicidae) as bio-indicators. *Biodiv. Conserv.*, 8, 1683–1695.

Ulloa-Chacon, P. & Cherix, D. (1994). Perspectives on control of the little fire ant (*Wasmannia auropunctata*) on the Galapagos Islands. In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 63–72). Boulder CO: Westview Press.

Vail, K. M. & Williams, D. F. (1994). Foraging of the Pharaoh ant, *Monomorium pharaonis*: an exotic in the urban environment. In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 228–239). Boulder CO: Westview Press.

van Pelt, A. F. (1966). Activity and density of old-field ants of the savannah river plant, South Carolina. *J. Elisha Mitchell Sci. Soc.*, 82, 35–43.

Vandervoude, C., Andersen, A. N. & House, A. P. N. (1997). Community organisation, biogeography and seasonality of ants in an open forest of south-eastern Queensland. *Aust. J. Zool.*, 45, 523–537.

Vanderwoude, C., Andersen, A. N. & House, A. P. N. (1997). Ant communities as bio-indicators in relation to fire management of spotted gum (*Eucalyptus maculata* Hook.) forests in South-East Queensland. *Mem. Mus. Vict.*, 56, 671–675.

Vanderwoude, C., Lobry de Bruyn, A., House, A. P. N. (2000). Response of an open-forest ant community to invasion by the introduced ant, *Pheidole megacephala*. *Austral Ecol.*, 25, 253–259.

Vanderwoude, C., Lobry de Bruyn, A. & House, A. P. N. (2000). Long-term ant community responses to selective harvesting of timber from spotted gum (*Corymbia variegata*)-dominated forests in south-east Queensland. *Ecol. Manag. Restor.*, 1, 204–214.

Vanderwoude, C. & Lobry de Bruyn, A. (2000). Short-term responses by ants (Hymenoptera: Formicidae) to selective timber harvesting of an open forest in eastern Australia. *Aust. For.*, 63, 185–194.

Vepsäläinen, K. & Wuorenrinne, H. (1978). Ecological effects of urbanization on the mound-building *Formica* L. species. *Mem. Zool.*, 29, 191–202.

Vepsäläinen, K., Savolainen, R., Tiainen, J. & Vilén, J. (2000). Successional changes of ant assemblages: from virgin and ditched bogs to forests. *Ann. Zool. Fennici*, 37, 135–149.

Vernier, R. (1995). Recherche écofaunistique sur les fourmis du genre *Formica* L. de la Tourbière du Cachot (Jura Neuchatelois) et Hauts-Marais Voisins (Hymenoptera, Formicidae) I. Liste des espèces et leurs biotopes préférentiels. *Bull. Soc. Neuchatel Sci. Nat.*, 118, 61–82.

Völkl, W., Woodring, J., Fischer, M., Lorenz, M. W. & Hoffmann, K. H. (1999). Ant-aphid mutualisms: the impact of honeydew production and honeydew sugar composition on ant preferences. *Oecologia*, 118, 483–491.

Wagner, D. & Gordon, D. M. (1999). Colony age, neighborhood density and reproductive

- potential in harvester ants. *Oecologia*, 119, 175–182.
- Waloff, N. & Blackith, R. E. (1962). The growth and distribution of the mounds of *Lasius flavus* (Fabricius) (Hym: Formicidae) in Silwood Park, Berkshire. *J. Anim. Ecol.*, 31, 421–437.
- Wang, C., Strazanac, S. & Butler, L. (2000). Abundance, diversity, and activity of ants (Hymenoptera: Formicidae) in oak-dominated mixed Appalachian forests treated with microbial pesticides. *Environ. Entomol.*, 29, 579–586.
- Wang, C., Strazanac, J. & Butler, L. (2001). A comparison of pitfall traps with bait traps for studying leaf litter ant communities. *J. Econ. Entomol.*, 94, 761–765.
- Warburg, I. & Steinberger, Y. (1997). On the spatial distribution of nests of the ants *Messor arenarius* and *Messor ebeinus*. *J. Arid Environ.*, 36, 671–676.
- Ward, P. S. (1987). Distribution of the introduced Argentine Ant (*Iridomyrmex humilis*) in natural habitats of the Lower Sacramento Valley and its effects on the indigenous ant fauna. *Hilgardia*, 55, 1–16.
- Way, M. J., Cammell, M. E., Paiva, M. R. & Collingwood, C. A. (1997). Distribution and dynamics of the Argentine ant *Linepithema (Iridomyrmex) humile* (Mayr) in relation to vegetation, soil conditions, topography and native competitor ants in Portugal. *Insect. Soc.*, 44, 415–433.
- Weber, S. (2002). Untersuchungen zur Ameisenfauna (Hymenoptera, Formicidae) einer Wildflusslandschaft (Tennengau, Salzburg). *Entomologica Austriaca*, 5, 16–17.
- Weber, S. (2003). Faunistisch-ökologische Untersuchungen der Ameisenfauna (Hymenoptera: Formicidae) einer Wildflusslandschaft im Salzburger Tennengau. *Myrmecologische Nachrichten*, 5, 15–30.
- Wesenigk-Sturm, B. (2002). Fundortbeschreibungen der Kerbameise *Formica (Coptoformica)*

foreli Emery 1909. *Ameisenschutz aktuell*, 16, 65–75.

Wesselinoff, G. (1979). Verbreitung und Schutz der Waldameisen in Bulgarien. *Bull. SROP*, 2, 497–498.

Wetterer, J. K. (1998). Nonindigenous ants associated with geothermal and human disturbances in Hawai'i Volcanoes National Park. *Pac. Sci.*, 52, 40–50.

Wetterer, J. K., Banko, P. C., Laniawe, L. P., Slotterback, J. W. & Brenner, G. J. (1998). Nonindigenous ants at high elevations on Mauna Kea, Hawai'i. *Pac. Sci.*, 52, 228–236.

Wetterer, J. K., Miller, S. E., Wheeler, D. E., Olson, C. A., Polhemus, D. A., Pitts, M., Ashton, I. W., Himler, A. G., Yospin, M. M., Helms, K. R., Harken, E. L., Gallaher, J., Dunning, C. E., Nelson, M., Litsinger, J., Southern, A. & Burgess, T. L. (1999). Ecological dominance by *Paratrechina longicornis* (Hymenoptera: Formicidae), an invasive tramp ant, in Biosphere 2. *Florida Entomol.*, 82, 381–388.

Whitford, W. G., van Zee, J., Nash, M. S., Smith, W. E. & Herrick, J. E. (1999). Ants as indicators of exposure to environmental stressors in North American desert grasslands. *Environ. Monit. Assess.*, 54, 143–171.

Wisniewski, J. & Dudek, J. (1974). Änderungen der Nestanzahl der Ameisen aus der *Formica rufa* - Gruppe in der Oberförsterei Proszkow während einem Zeitablauf von 5 Jahren. *Pr. Kom. Nauk Roln. Kom. Nauk Lesn. Poznan. Tow. Przyj. Nauk Wydz. Nauk Roln.*, 38, 239–247.

Wojcik, D. P. (1994). Impact of the red imported fire ant on native ant species in Florida. In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 269–281). Boulder CO: Westview Press.

Woyciechowski, M. & Miszta, A. (1976). Spatial and seasonal structure of ant communities in a mountain meadow. *Ekol. Polsk.*, 24, 577–592.

- Wojciechowski, M. (1985). The ants (Hymenoptera, Formicidae) of the Male Pieniny Mountains - Carpathians. *Acta Zool. Cracov.*, 28, 283–296.
- Wojciechowski, M. (1993). Ants (Hymenoptera, Formicidae) of the glades in the Tatra Mts (The Carpathians). *Tiscia*, 27, 17–22.
- Wright, P. J. (1989). Energy turnover in a population of the ant *Lasius flavus* F. *Actes coll. Insect. Soc.*, 5, 69–73.
- Wuorenrinne, H. (1983). Der Einfluß der Waldsplitterung auf Ameisengesellschaften am Beispiel von Espoo (Finnland). *Tagungsber 2. Leipzig. Symp. urb. Ökol.* 1983, 25–30.
- Wuorenrinne, H. (1989). Effects of urban pressure on colonies of *Formica rufa* group (Hymenoptera. Formicidae) in the town of Espoo (Finland). *Ann. Zool. Warsz.*, 42, 335–343.
- Xu, Z., Hu, G. & Yu, X. (1999). Biomass and ecological function of ant communities in the tropical rain forest of Xishuangbanna, China. *Zool. Res.*, 20, 441–445.
- York, A. (1994). The long-term effects of fire on forest ant communities: management implications for the conservation of biodiversity. *Mem. Queensl. Mus.*, 36, 231–239.
- York, A. (1999). Long-term effects of frequent low-intensity burning on the abundance of litter-dwelling invertebrates in coastal blackbutt forests of southeastern Australia. *J. Insect Conserv.*, 3, 191–199.
- York, A. (2000). Long-term effect of frequent low-intensity burning on ant communities in coastal blackbutt forests of southeastern Australia. *Austral Ecol.*, 25, 83–98.
- Zakharov, A. A. (1990). Vertical structure of the ant communities and their role in the Peruvian selva. *Mem. zool.*, 44, 7–11.
- Zenner-Polania, I. (1994). Impact of *Paratrechina fulva* on other ant species. In Williams, D. F. (Ed.), *Exotic ants. Biology, impact, and control of introduced species* (pp. 121–132). Boulder CO: Westview Press.

Zorrilla, J. M., Serrano, J. M., Casado, M. A., Acosta, F. J. & Pineda, F. D. (1986). Structural characteristics of an ant community during succession. *Oikos*, 47, 346–354.